



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A SUPPOSED NEW SPECIES OF HUMMINGBIRD IN
THE ROYAL ZOOLOGICAL MUSEUM OF
DRESDEN.

BY A. B. MEYER.

Eriocnemis aurea sp. nov.

E. cupreiventri (Fras.) similis, sed minor, corpore toto, gula et pectore exceptis, aureo-æneo, caudæ tectricibus longis prasinis, subcaudalibus cobaltinis plus minusve viridi marginatis, tibiarum pappis minoribus, diversus.

Long. al. 58; caud. 40; rostri 20 mm.

HAB. — Colombia.

E. cupreiventris measures: wing, 62 mm.; tail, 45 mm., bill of equal length. The coloration of the specimen above described differs so considerably from *cupreiventris* that it cannot be taken for an individual variation. While *cupreiventris* is only slightly tinted with bronze, *aurea* is covered with golden bronze all over, and in part of a most lively tint, even on the shorter tail and wing coverts. The throat and upper breast are lighter green than in *cupreiventris*, and in certain lights washed with bronze too, and each feather has a concealed white cross band as in *E. dybowskii* Tacz. Further, the long upper tail-coverts are not greenish blue, but green, and the under tail-coverts are not violet, but cobalt blue, only in certain lights they are hyacinth-blue, some margined with green. The color of the wings as well as of the tail is not as lively as in *cupreiventris*; and, finally, the white plumes on the thighs are much shorter and not as abundant. The specimen cannot be taken for a young *cupreiventris*, being much more brightly colored on the whole body, than the adult of this species.

SOMATERIA DRESSERI, THE AMERICAN EIDER.

BY GEORGE H. MACKAY.

THIS bird is found, as far as I have been able to learn, only on the Atlantic coast from Labrador to Delaware, collecting in large